

# Density dependence in social behaviour: home range overlap and density interacts to affect conspecific encounter rates in a gregarious ungulate

Eric Vander Wal · Michel P. Laforge ·  
Philip D. McLoughlin

Received: 4 March 2013 / Revised: 5 November 2013 / Accepted: 5 November 2013  
© Springer-Verlag Berlin Heidelberg 2013

**Abstract** Sociality is poorly understood in the context of population processes. We used wild, female elk (*Cervus canadensis*) equipped with proximity-logging radio collars ( $n=62$ ) from Manitoba, Canada (2007–2009), to test for modifying effects of population density (two areas: 0.42 and 0.22 animals/km<sup>2</sup>) on the relationship between two measures of sociality. This included the rate at which collared individuals encountered one another per year (encounters logged as animals ranging to within 1.4 m of each other) and the extent to which animals overlapped in annual home range (proportion of shared minimum convex polygon ranges). Overlap was significantly greater in the high density area compared to that of the low, but not if we only considered individuals that directly encountered each other, implying that familiar individuals will maintain a constant degree of range overlap regardless of density. Encounter rate was nonlinearly related to home range overlap. This relationship was also density-dependent, exhibiting negative density dependence at high proportions of overlap, primarily in the high density subpopulation. Sociality, as defined by two interacting measures of behaviour—encounter rate and home range overlap—exhibits a complex nonlinear relationship; we discuss the implications

of these results as they pertain to sociobiology, resource competition, and pathogen transmission.

**Keywords** *Cervus canadensis* · Density dependence · Elk · Encounter rate · Sociality

## Introduction

Sociality is broadly defined as the properties and processes of social existence (Wilson 1975). Ecologists interested in sociality seek to inform us about the ecological and evolutionary factors that promote and maintain interactions between conspecifics leading to group living. Despite broad acceptance that density-dependent processes are fundamental to ecology, density dependence in the study of sociality has thus far received limited attention (but see Bateman et al. 2012; Vander Wal et al. 2012a, 2013a; Cross et al. 2013).

Sociality in group-living species can be quantified through multiple measures, for example, when two individuals directly encounter each other and express social behaviours (i.e., dyadic interactions or encounters). Dyadic encounters provide the foundation for analysing social structure, as interactions form the basis for relationships between individuals (Hinde 1976; Whitehead 1997). From dyadic interactions, a second level of organisation is often assumed to emerge. This occurs at the extent where individuals are observed to share the same area or range. For example, associations of larger movement patterns over longer time periods that reflect group associations and shared (or avoided) space use. The reverse assumption—that such spatial patterns imply fine-scale interactions—may be referred to as the ‘gambit of the group’ (Franks et al. 2010). Here, individuals seen grouping together (e.g., sharing home ranges or travelling together in a school or herd) are assumed to be directly interacting with every other individual in that group.

---

Communicated by K. E. Ruckstuhl

**Electronic supplementary material** The online version of this article (doi:10.1007/s00265-013-1652-0) contains supplementary material, which is available to authorized users.

---

E. Vander Wal · M. P. Laforge · P. D. McLoughlin  
Department of Biology, University of Saskatchewan, 112 Science  
Place, Saskatoon SK S7N 5E2, Canada

*Present Address:*

E. Vander Wal (✉)  
Département de biologie, Université de Sherbrooke, 2500 boul. de  
l’université, Sherbrooke QC J1K 2R1, Canada  
e-mail: Eric.Vander.Wal@USherbrooke.ca

But do high levels of association in space always imply high rates of encounter between animals (dyads)? If a group is formed due to the fusion of two smaller groups, individuals of founding groups are known to interact with one another at different rates after the fusion event (Millsbaugh et al. 2004a; Couzin 2006). Also, familial associations (Dubost 1970; Clutton-Brock et al. 1982; Wronski and Apio 2006) may bias the relationship between large-scale overlap in movements and fine-scale interactions. These are considerations internal to the dynamics of a group, but how well spatial overlap and indirect associations match with direct encounters of dyads might also be affected by the extrinsic ecological factors that define the costs and benefits of group living.

Foremost among external factors affecting group dynamics is conspecific density and the process of resource competition, e.g., for food or mates. Density is known to affect group size and composition (Ruckstuhl and Festa-Bianchet 2001; Pépin and Gerard 2008; Vander Wal et al. 2013a). It has further been shown to affect dyadic encounter rates and encounter duration in experimental settings (Vander Wal et al. 2012a), and to increase agonistic behaviour (Pettit-Riley et al. 2002; Moyle et al. 2009). Density is also known to decrease mobility (e.g., ‘social fence’ hypothesis (Matthysen 2005)), necessitating an increase in home range overlap with higher abundance (e.g., rodents (Ostfeld et al. 1985; Ferron and Ouellet 1989; Erlinge et al. 1990; Steury and Murray 2003), felids (Nielsen and Woolf 2001), and ursids (Dahle and Swenson 2003)). Conversely, very low population densities may have consequences to social behaviour and breeding ecology, giving rise to Allee effects (Brashares et al. 2010). If densities directly follow from resource availability (i.e., habitat *sensu* Morris (2003)), nonlinearity in range separation (territoriality) may be apparent, with overlap occurring at low and very high levels of density (food abundance (see White 2004 for review)), but not moderate levels where the relative costs of food or territory defence are low. This situation has been observed in bears (McLoughlin et al. 2000), bats (Chaverri et al. 2007), and birds (Carpenter and Macmillan 1976). Recognising the complex nature of how home range overlap is determined and how density might affect encounter rates, what can we expect for the relationship between range overlap and the rate of direct encounter between dyads as density increases? This aspect of sociality is poorly understood.

Quantifying the relationship between home range overlap and encounter rate is of vital importance to studies on the evolution of sociality and to assess the risk of pathogen transmission, especially in studies where home range overlap is used as a proxy for encounter (or even contact) rates (see Whitehead and Dufault (1999) for review; Robert et al. 2012). Here, we test the prediction of increased encounter rate with increasing home range overlap. From this prediction, we constructed a hierarchical set of competing models (Tables 1 and S1). These include that patterns in encounter rate can be

**Table 1** *A priori* models and model selection for dyadic encounter rate per year (response variable) as a function of proportion of home range overlap (HRO) for female elk in two subpopulations of Riding Mountain National Park, Manitoba, Canada (2007–2009): the east low density subpopulation (0.22 elk/km<sup>2</sup>) and the west high density subpopulation (0.42 elk/km<sup>2</sup>). Models compose three concepts of increasing complexity: (1) simple linear relationships ( $M_1$ ,  $M_2$ ) or additive effects ( $M_3$ ) can describe the relationship between HRO, density, and encounter rate; (2) a feedback exists between HRO and density ( $M_4$ ); and (3) nonlinearity in encounter rate by HRO results from possible competitive effects or avoidance ( $M_5$ ); or the global model ( $M_6$ ). See Supplementary Table S1 for further details

Models	Independent	$\Delta AIC^a$	$AIC_w^b$
$M_1$	HRO	>10	0
$M_2$	Density	>10	0
$M_3$	HRO+Density	>10	0
$M_4$	HRO+Density+HRO*Density	>10	0
$M_5$	HRO+HRO <sup>2</sup> +Density	0	0.61
$M_6$	HRO+HRO <sup>2</sup> +Density+HRO*Density	0.9	0.39

<sup>a</sup> Change in Akaike’s Information Criteria value

<sup>b</sup> Akaike’s Information Criteria model weight

explained by home range overlap and subpopulation density independently ( $M_1$ ,  $M_2$ ) or their additive effect ( $M_3$ ). Alternately, home-range overlap interacts with density to create a positive feedback ( $M_4$ ) or, possibly due to competitive effects, nonlinearity exists between home range overlap and encounter rate ( $M_5$ ). Finally, we evaluate a global model encompassing both the home range overlap by density feedback and the nonlinearity predictions ( $M_6$ ). We test these models using wild, female elk ( $n=62$ ) equipped with proximity-logging radio collars from two discrete subpopulations in Riding Mountain National Park (RMNP), Manitoba, Canada (2007–2009), which vary markedly in their density. Female elk are highly gregarious, and we assumed that the higher the overlap in home range, the more strongly individuals were associated with the group using that range (Franklin et al. 1975; Millsbaugh et al. 2004a). We discuss the implications of our findings as it relates to the density dependence of sociality and the applications to sociobiology and modelling the dynamics of communicable disease (e.g., in the elk of RMNP which are affected by bovine tuberculosis, TB).

## Methods

### Study site and animals

Riding Mountain National Park, Manitoba, Canada (RMNP, 3,000 km<sup>2</sup>, Fig. S1) lies on the ecotone of the Boreal Plains and Prairie Parkland biomes (Rowe 1972; Caners and Kenkel 2003). An elevation gradient exists in RMNP: the eastern portions of the park rise 475 m from the Manitoba lowlands

up the Manitoba escarpment. As a result, variation exists in the habitat. At high elevations, the boreal forest dominated by mixedwood forests includes spruce (*Picea Marianna* and *P. glauca*), pine (*Pinus banksiana*), and aspen (*Populus tremuloides*) that are interspersed with waterways, open fens, and bogs. At lower elevations, open marshes and remnant fescue prairie are interspersed in a matrix of aspen parkland and mixedwood forests. RMNP is essentially a closed system surrounded by an agricultural matrix (for further details, see Vander Wal 2011).

RMNP is a refuge for many species of wildlife, including a population of free-ranging elk. There are two distinct subpopulations of elk in the park (Vander Wal et al. 2012b, 2013b) which occur at different densities: east 0.22 elk/km<sup>2</sup> and west 0.42 elk/km<sup>2</sup> (hereafter ‘low’ and ‘high’ relative density, respectively). Subpopulation densities were derived from a 3-year average (2007–2009) of a 25 % coverage aerial survey of visible elk (Parks Canada unpublished data, see Vander Wal 2011 and Vander Wal et al. 2013a for details of survey design).

Elk are considered the reservoir host for bovine tuberculosis (TB, *Mycobacterium bovis*) in the region (see Brook and McLachlan (2006) and Shury and Bergeson (2011) for details). Presence of TB has had considerable economic impacts on the region (Nishi et al. 2006; Brook and McLachlan 2006) as elk are known to co-mingle with cattle (*Bos taurus*; Brook et al. 2013). TB is, however, predominantly restricted to the western subpopulation of RMNP elk (Shury and Bergeson 2011; Vander Wal et al. 2012b).

#### Proximity-logging radio collars

Proximity-logging radio collars (SirTrack, Havelock North, New Zealand) are a recent technology which quantify encounters between two marked individuals whenever they are within a pre-determined distance from one another (see Cross et al. 2012 for review). Measures of spatial proximity (hereafter encounters) between individuals are used to infer social behaviours such as interactions and direct contact between individuals (e.g., see Ji et al. 2005; Prange et al. 2006; Walrath et al. 2011; Cross et al. 2012; Robert et al. 2012, 2013; Vander Wal et al. 2012a, b; Vander Wal et al. 2013c for examples). Traditionally, studies have relied on radio-tracking data to infer dyadic encounters between individuals, with an encounter (typically referred to as a ‘contact’) being defined as broadly as two individuals logging spatial locations within 100 m of each other (White and Harris 1994; Ramsey et al. 2002; Schaubert et al. 2007; Kjaer et al. 2008). The advent of proximity collars has greatly improved our ability to continuously detect close spatial associations (Ji et al. 2005; Prange et al. 2006; Goodman 2007; Cross et al. 2012) at scales, in locations, and at times previously unavailable and inaccessible with either traditional ethological techniques or satellite- and radio-telemetry.

We programmed proximity collars to record dyadic encounters whenever two collared elk came to within  $1.42 \pm 0.20$  m ( $\bar{x} \pm SD$ ) of one another, (Goodman 2007). For technical reasons (see Goodman 2007), collars deactivated recording an encounter at a distance of  $1.98 \pm 1.60$  m each time a pair separated for  $>30$  s ( $\bar{x} \pm SD$ , Goodman 2007). As individuals were not observed directly interacting, or physically contacting, we are left to assume that these social behaviours (i.e., indicators of sociality) were being exhibited when the two animals are engaged in an encounter with one another; we assume these encounters do not occur randomly due to the highly social nature of elk. For encounters to be recorded, the elk needed to be within one body length of a collared conspecific. For further discussion regarding proximity collars and data treatment, see Robert et al. (2013) and Vander Wal et al. (2013c).

#### Animals

We used a net-gun fired from a helicopter (Cattet et al. 2004) to capture female elk ( $n=24$  in 2007 and  $n=15$  in 2008 [high density subpopulation] and  $n=23$  in 2008 [low density subpopulation]). We equipped each animal with proximity collars. Collars were deployed simultaneously and recorded encounters for all dyads within the replicate for 1 year (February–February). The female elk in our study represented a random sample of the adult population, and no correlation between spatial proximity and genetic relatedness was found (Vander Wal et al. 2012b). We relocated the animals several times throughout the year ( $\bar{x} = 30.4$ ) by fixed-wing aircraft (Cessna 172) during daylight hours, usually 0800–1900 h, two to four times per month (see Vander Wal et al. 2011 for details). Animals were collared for equivalent time and collars either dropped off via mechanical release or animals were recaptured and collars removed.

#### Home range overlap

We determined home range overlap of tracked elk using ArcMap (ArcGIS 9.3, ESRI Redlands California, USA) and the Home Range Tools extension (Rodgers et al. 2005) to calculate 95 % minimum convex polygons. Minimum convex polygons were used due to their ease in comparison across studies and a limited number of relocations (see Fig. S2 for further discussion). We used an index of overlap based on the formula presented by (Millsbaugh et al. 2004b):

$$\text{Overlap} = \frac{A_{1,2}}{(A_1 + A_2) - A_{1,2}} \quad (1)$$

where  $A_1$  and  $A_2$  are the areas of the home ranges of animals 1 and 2, respectively, and  $A_{1,2}$  is the area of overlap between

individual 1 and 2. (See Fig. S3 for an example of low, medium, and high range overlap).

### Statistical analysis

We tested the relationship between dyadic encounter rate among dyads and home range overlap at the two subpopulation densities using six *a priori* models (Table 1). The simplest of which predicts that univariate tests of whether home range overlap or subpopulation density independently best predict encounter rate ( $M_1$ ,  $M_2$ ). Subsequently, we test whether subpopulation density and home range overlap ( $M_3$ ) and the interaction between the two ( $M_4$ ) better predict encounter rate. Lastly, we test whether these are better described by including a quadratic term to describe the non-linearity in the relationship ( $M_5$ ,  $M_6$ ). Models were ranked based on their parsimony evaluated using Akaike's Information Criteria (AIC; Burnham and Anderson 2002). All models with a  $\Delta AIC < 2$  were considered indistinguishable. Indistinguishable models were consequently averaged using the R package MuMIn (Barton 2010).

Frequencies of encounters were count data; therefore, all generalised linear models included a Poisson distributed error term. Because logically an animal with no overlap in their home ranges cannot have had fine-scale pairwise encounters, we forced the regression through the origin. We added subpopulation density to the model as a fixed two-level factor ( $M_{2-6}$ ) and its interaction with home range overlap ( $M_{4 \text{ and } 6}$ ). Our dyads were not replicated across years, precluding the use of mixed-effects generalised linear models typically used to control pseudo-replication in pairwise observations. Instead, we randomised the generalised linear model (Manly 1998) to generate a distribution of  $z$ -values against which we tested the significance of each covariate. We ran 1,000 iterations of the linear regression with  $n=1,000$  randomised samples. This does not fully account for pseudo-replication, but it does ensure that patterns in these data are not statistical artefacts. All statistics were performed in R version 2.15 (2012).

### Results

The mean number of encounters per dyad was  $65.7 \pm 9.2$  (both densities pooled). Dyads at high ( $0.42 \text{ elk/km}^2$ ) density had more encounters ( $83.8 \pm 12.4$ ,  $n=709$ ) than dyads at low ( $0.22 \text{ elk/km}^2$ ) density ( $16.7 \pm 4.4$ ,  $n=506$ ).


The mean home range size of female elk was  $42.5 \pm 4.5 \text{ km}^2$  ( $\bar{x} \pm \text{SE}$ ) in the high density subpopulation and  $34.4 \pm 5.9 \text{ km}^2$  in the low density subpopulation. Mean home range overlap per dyad among all dyadic pairs in each study site varied between study areas, with overlap greater in the higher density subpopulation ( $0.13 \pm 0.007$ ) than in the lower density subpopulation ( $0.07 \pm 0.006$ ). However, when non-interacting dyads were excluded from the analysis of home range overlap,

overlap was not different in the two density treatments (high:  $0.18 \pm 0.008$ , low:  $0.18 \pm 0.012$ ; Fig. 1).


Frequency of encounters was nonlinearly related to home range overlap and varied by subpopulation density (Fig. 2). *A priori* models  $M_5$  and  $M_6$ , including subpopulation density, home range overlap, its quadratic term ( $M_5$ ), and the interaction between home range overlap and subpopulation density ( $M_6$ ), were the most parsimonious ( $AIC_w$ ,  $M_5=0.61$ ,  $M_6=0.39$ ; Table 1). This was despite the additional interaction term in  $H_6$  being nonsignificant ( $z=-1.05$ ,  $p=0.29$ ; Table 2). Home range overlap and subpopulation density all had significant effects in the models ( $p < 0.001$ ; Table 2). Results did not differ for models where the intercept was permitted to vary from 0 (data not shown).

### Discussion

Density dependence in social behaviour can have important implications for population dynamics (Bateman et al. 2012) and disease prevalence (Lloyd-Smith et al. 2005). In both instances, the mechanisms of these density-dependent effects occur at the fine spatial scale of pairwise encounters, e.g., interference competition or pathogen transmission (Krause and Ruxton 2002). However, the effect of density on these encounters has rarely been explored (but see Vander Wal et al. 2012a; Cross et al. 2013). Here, we described not only density dependence but also how it is modified in a nonlinear fashion



**Fig. 1** Mean proportion of home range overlap in female elk of Riding Mountain National Park, Manitoba, Canada (2007–2009). **a** Represents the mean (95 % CI) when all dyads were considered within each study region; **b** the mean proportion of shared space when only individuals with overlapping ranges were considered. Overlapping dyads shared similar home range overlap at both densities, but overlapped with more individuals at higher densities



**Fig. 2** Encounter rate per dyad versus home range overlap for female elk in Riding Mountain National Park, Manitoba, Canada (2007–2009) for two subpopulations experiencing different densities: high density (0.42 elk/km<sup>2</sup>, triangles); and low density, 0.22 elk/km<sup>2</sup> (grey circles). Predicted values from a generalised linear model (averaged  $M_5$  and  $M_6$ , see Table 2) are plotted as lines with 95 % CI (high density in black, low density in grey). Inset, predictions plotted in anti-log space, illustrate how encounter rates saturate (for low density, grey line) and exhibit negative density dependence (at high density, black line)

by shared space use between individuals. For example, due to nonlinear effects thresholds appear where increased home range overlap may lead to a saturated frequency of fine-scale dyadic encounters (Fig. 1, low density subpopulation) or even avoidance of fine-scale dyadic encounters (Fig. 1, high density subpopulation). Saturation of encounters and avoidance

**Table 2** Averaged ( $M_5$ ,  $M_6$  Table 1) generalised linear model with Poisson error distribution for dyadic encounter rate per year by proportion of home range overlap for female elk in two subpopulations of Riding Mountain National Park, Manitoba, Canada (2007–2009): the east, low density (0.22 elk/km<sup>2</sup>) and the west, high density (0.42 elk/km<sup>2</sup>)

	Coefficient (Unconditional SE)	z-value <sup>a</sup>	P-value
HRO <sup>b</sup>	18.04 (0.13)	135.05 (−40.93–46.24)	<0.001
HRO <sup>2</sup>	−17.18 (0.13)	130.10 (−48.00–46.77)	<0.001
Low density	−0.55 (0.04)	−13.62 (331.42–557.47)	<0.001
High density	1.17 (0.01)	60.00 (327.13–531.61)	<0.001
HRO* density	−0.14 (0.13)	−1.05 (−51.63–43.03)	0.294

<sup>a</sup> 95 %CI generated by randomising the frequency of encounters in a generalised linear model to test whether the outcome could have been generated randomly and thus may have been a function of pseudo-replication due to pairwise data

<sup>b</sup> Home range overlap

suggests that benefits of social behaviour may become costs when two individuals share a large extent of their home range.

Within the shared space, individuals compete over resources (Rieucan and Giraldeau 2011). Direct competition may require more encounters at higher density than low, regardless of how much range is actually shared between individuals. Weckerly (1999) found that aggressive interactions increased in elk with increasing population density, suggesting that as resources became scarcer, defending them was more important. Conversely, in an experimental manipulation of domestic sheep, Mobæk et al. (2012) found there was no evidence for interference competition or the defence of resources. They did, however, demonstrate that animals at high density (low resource abundance) spent more time active. Being active increases the probability of encountering conspecifics. Indeed, Vander Wal et al. (2012a) found that increasing density increased encounter rates in an experimental setting that was not resource-limited. One hypothesis which follows is that increased encounter rates may be proximally attributed to resource availability, but ultimately a function of social phenomena, such as allogrooming or maintaining a hierarchy.

Nevertheless, the nonlinearity and density dependence of our results imply an important, yet unexplored, interplay between local habitat (e.g., structure, resource availability) and encounter rate—one that might vary geographically across the range of our study species. For example, in ungulates open habitats—such as those found in RMNP—are known to result in larger groupings due to emergent group properties (Gerard and Loisel 1995) or increased predation risk (Hebblewhite and Pletscher 2002). Increased group size has been shown to increase encounter rate (Cross et al. 2013). Increased gregariness also precipitates increased home range overlap. Conversely, complex habitats break groups apart (Haydon et al. 2008; Fortin et al. 2009), decreasing the shared space use between individuals. As a result, open habitats are predicted to increase home range overlap and encounter rates. For elk, open habitats with abundant graminoids are known to be preferred habitats (Christianson and Creel 2007), likely supporting higher densities of animals.

Higher densities of elk also result in larger groups (Vander Wal et al. 2013b). Larger groups may be composed of more familiar individuals, as individuals in larger groups may have been able to associate on a smaller scale with more individuals over longer periods of time than those living at lower densities (smaller groups). Familiar animals will interact with one another at higher rates (Millspaugh et al. 2004a; Couzin 2006). Albeit these individuals are often kin; here, encounters among elk are equally likely to be among unrelated individuals (Vander Wal et al. 2012b). Nevertheless, it may be that females must experience a trade-off between interference competition (direct encounters) with other individuals versus scramble competition (avoiding or sharing less space with

conspecifics and lower home range overlap). Or, similarly, the trade-off may occur along a gradient of conflict within interference competition (e.g., proximal avoidance versus avoiding physical contact). These trade-offs may be contingent on the relationships that exist between animals at different densities, the availability of resources within home ranges, and their seasonal use (see Vander Wal et al. 2013c for details on seasonal encounter rates).

Albeit infrequently tested (but see Robert et al. 2012), it has been assumed that home range overlap can be used as a proxy for encounter rate (see review by Whitehead and Dufault (1999)). Certainly our findings corroborate this finding; however, with the important caveat that the relationships can be expected to be both nonlinear and density-dependent. Therefore, this assumption may be flawed due to the fact that it is likely site-specific. One area where this assumption has been applied is in wildlife disease ecology.

Sociobiology and disease ecology are quickly finding common ground, as the mechanisms for horizontal pathogen transmission involve individual social encounters (Tompkins et al. 2011). Home range overlap has been used as a proxy for risk of disease transmission (Caley et al. 1998; Ramsey et al. 2002; Schaubert et al. 2007). Although our results confirm a correlation between home range overlap and encounter rate, they also caution that the relationship, similar to disease transmission, can be modified by host density. Our results indicate that at high population density, female elk compensate by sharing space with more conspecifics, but share the same amount of space per dyad than at lower density. This may result in increased disease transmission as infected hosts will associate more with susceptible conspecifics, an important element in disease transmission (McCallum et al. 2001). However, as home range overlap increases encounter rates saturate and are predicted to decrease, suggesting that elk are avoiding fine-scale social interactions when shared space use is high (e.g., >40 %). This result varies in magnitude. Nevertheless, it illustrates disproportionately fewer encounters, which therefore may predict decreased pathogen transmission. Some social behaviours, e.g., group size, have been shown to increase the prevalence of directly transmissible pathogens and parasites (Côté and Poulin 1995). Fine-scale social avoidance, however, could be a mechanism to mitigate this transmission. This relationship, however, is also affected by conspecific density. For example, increased density has been shown to increase the risk of parasitic infection, which consequently affects annual reproductive success (Albon et al. 2002).

Our study was a large-scale natural ‘experiment’ which capitalised on existing natural variation between discrete subpopulations. A strength of our design is that it occurred in a wild population (cf., captive elk used in Vander Wal et al. 2012a). A weakness, however, was failure to replicate density. Although, the relationship between home range overlap and encounter rate was well-replicated, our density treatment was perfectly pseudo-replicated (Hurlbert 1984)—technically

limiting our capacity to make inferences outside of our study system. Given that density has been shown to affect encounter rates (Vander Wal et al. 2012a) and home range overlap (e.g., Steury and Murray 2003; Nielsen and Woolf 2001; Dahle and Swenson 2003), we submit that density likely had a modifying effect on the relationship between home range overlap and encounter rate in our study. However, density itself may be affected by other differences between subpopulations, including habitat, which we do not disentangle here. Indeed, due to density-dependent habitat selection theory, some authors (e.g., Morris 2003) suggest that the definition of habitat needs to include both resources and density.

In summary, our results identified three key findings. First, encounter rate is affected by shared space use; thus, illustrating a possible mechanism to minimise competition and pathogen transmission by avoiding fine-scale encounters when shared space use is high. Second, this relationship is density-dependent, with encounters occurring at dramatically higher rates in high density landscapes. Ultimately, this questions the validity of assuming that shared space use correlates directly with encounter rate. Although a correlation clearly exists, it appears to be context-dependent; as such, we caution against generalising between systems. In addition to these findings, we have identified two important gaps in research into density-dependent social behaviour. The first is to better understand the role of habitat (e.g., structure and resource availability) in shaping social encounters. The second is to highlight the need to integrate data derived from proximity-detecting technology with traditional ethological techniques to identify which social behaviours are being expressed within each encounter.

**Acknowledgments** Funding and logistical support was provided primarily by Parks Canada Agency, the Natural Science and Engineering Research Council of Canada (F. Messier, P.D. McLoughlin), and PrioNet Canada. P. Simpson and B. Simpson adroitly flew elk relocation flights and C. Wilson, T. Vandenbrink, and T. Shury efficiently and safely handled elk. R. Grzela, S. McKay, R. Robinson, S. Johnstone, J. Dillabough, B. Blackbird, S. Boychuk, S. Helms, A. Ledden, and M. Benson assisted with monitoring collared animals. Thanks to T. Sallows, D. Bergeson, and K. Kingdon at Parks Canada for their commitment to this research and to F. Messier, R. K. Brook, and F. Pelletier for support while writing this manuscript. Also, thanks to Kathreen Ruckstuhl and a number of anonymous reviewers whose suggestions markedly improved our work.

**Ethical Standards** Our study, which occurred in Canada, complied with the current laws of the country at the time. Our research followed Animal Care Protocol #20060067 of the University of Saskatchewan following the guidelines of the Canadian Council on Animal Care.

## References

- Albon SD, Stien A, Irvine RJ, Langvatn R, Ropstad E, Halvorsen O (2002) The role of parasites in the dynamics of a reindeer population. *Proc R Soc Lond B* 269:1625–1632

- Barton K. (2010) MuMin: multi-model inference version 0.13.1 <http://CRAN.R-project.org/package=Mumin>
- Bateman AW, Ozgul A, Coulson T, Clutton-Brock TH (2012) Density dependence in group dynamics of a highly social mongoose, *Suricata suricatta*. *J Anim Ecol* 81:628–639
- Brashares JS, Werner JR, Sinclair ARE (2010) Social “meltdown” in the demise of an island endemic: allee effects and the Vancouver Island marmot. *J Anim Ecol* 79:965–973
- Brook RK, McLachlan SM (2006) Factors influencing farmers’ concerns regarding bovine tuberculosis in wildlife and livestock around Riding Mountain National Park. *J Environ Manage* 80:156–166
- Brook RK, Vander Wal E, van Beest FM, McLachlan SM (2013) Use of cattle winter feeding areas by elk and white-tailed deer: implications for managing bovine tuberculosis transmission risk from the ground up. *Prev Vet Med* 108:137–147
- Burnham KP, Anderson DR (2002) Model Selection and Multimodel Inference: A Practical Information-Theoretic. Springer, New York
- Caley P, Spencer NJ, Cole RA, Efford MG (1998) The effect of manipulating population density on the probability of den-sharing among common brushtail possums, and the implications for transmission of bovine tuberculosis. *Wildlife Res* 25:383–392
- Caners RT, Kenkel NC (2003) Forest stand structure and dynamics at Riding Mountain National Park, Manitoba, Canada. *Community Ecol* 4:185–204
- Carpenter FL, Macmillan RE (1976) Threshold model of feeding territoriality and test with a Hawaiian honeycreeper. *Science* 194:639–642
- Cattet MRL, Caulkett NA, Wilson C, Vandenbrink T, Brook RK (2004) Intranasal administration of xylazine to reduce stress in elk captured by net gun. *J Wildlife Dis* 40:562–565
- Chaverri G, Gamba-Rios M, Kunz TH (2007) Range overlap and association patterns in the tent-making bat *Artibeus watsoni*. *Anim Behav* 73:157–164
- Christianson DA, Creel S (2007) A review of environmental factors affecting elk winter diets. *J Wildlife Manage* 71:164–176
- Clutton-Brock TH, Guinness FE, Albon SD (1982) Red Deer: Behavior and Ecology of Two Sexes. University of Chicago Press, Chicago
- Côté IM, Poulin R (1995) Parasitism and group size in social animals: a meta-analysis. *Behav Ecol* 6:159–165
- Couzin ID (2006) Behavioral ecology: social organization in fission–fusion societies. *Curr Biol* 16:R169–R171
- Cross PC, Creech TG, Ebinger MR, Heisey DM, Irvine KM, Creel S (2012) Wildlife contact analysis: emerging methods, questions, and challenges. *Behav Ecol Sociobiol* 66:1437–1447
- Cross PC, Creech TG, Ebinger MR, Manlove K, Irvine KM, Henningsen J, Rogerson JD, Scurlock BM, Creel S (2013) Female elk contacts are neither frequency nor density dependent. *Ecology* 94:2076–2086
- Dahle B, Swenson JE (2003) Home ranges in adult Scandinavian brown bears (*Ursus arctos*): effect of mass, sex, reproductive category, population density and habitat type. *J Zool* 260:329–335
- Dubost G (1970) L’organisation spatiale et sociale de *Muntiacus reevesi* Ogilby 1839 en semi-liberté. *Mammalia* 34:331–355
- Erlinge S, Hoogenboom I, Agrell J, Nelson J, Sandell M (1990) Density-related home-range size and overlap in adult field voles (*Microtus agrestis*) in southern Sweden. *J Mammal* 71:597–603
- Ferron J, Ouellet JP (1989) Temporal and intersexual variations in the use of space with regard to social organization in the woodchuck (*Marmota monax*). *Can J Zool* 67:1642–1649
- Fortin D, Fortin M-E, Beyer HL, Duchesne T, Courant S, Dancose K (2009) Group-size-mediated habitat selection and group fusion–fission dynamics of bison under predation risk. *Ecology* 90:2480–2490
- Franklin WL, Mossman AS, Dole M (1975) Social organization and home range of Roosevelt elk. *J Mammal* 56:102–118
- Franks DW, Ruxton GD, James R (2010) Sampling animal association networks with the gambit of the group. *Behav Ecol Sociobiol* 64:493–503
- Gerard J-F, Loisel P (1995) Spontaneous emergence of a relationship between habitat openness and mean group size and its possible evolutionary consequences in large herbivores. *J Theor Biol* 176:511–522
- Goodman EL (2007) Quantifying interactions in a high-density badger (*Meles meles*) population. Dissertation, University of York
- Haydon DT, Morales JM, Yott A, Jenkins DA, Rosatte R, Fryxell JM (2008) Socially informed random walks: incorporating group dynamics into models of population spread and growth. *Proc R Soc Lond B* 275:1101–1109
- Hebblewhite M, Pletscher DH (2002) Effects of elk group size on predation by wolves. *Can J Zool* 80:800–809
- Hinde RA (1976) Interactions, relationships and social structure. *Man* 11:1–17
- Hurlbert SH (1984) Pseudoreplication and the design of ecological field experiments. *Ecol Monogr* 54:187–211
- Ji W, White PCL, Clout MN (2005) Contact rates between possums revealed by proximity data loggers. *J Appl Ecol* 42:595–604
- Kjaer LJ, Schaub EM, Nielsen CK (2008) Spatial and temporal analysis of contact rates in female white-tailed deer. *J Wildlife Manage* 72:1819–1825
- Krause J, Ruxton G (2002) Living in Groups. Oxford University Press, Oxford
- Lloyd-Smith JO, Cross PC, Briggs CJ, Daugherty M, Getz WM, Latto J, Sanchez MS, Simth AB, Swei A (2005) Should we expect population thresholds for wildlife disease? *Trends Ecol Evol* 20:511–519
- Manly BFJ (1998) Randomization, Bootstrap and Monte Carlo Methods in Biology. Chapman and Hall, London
- Matthysen E (2005) Density-dependent dispersal in birds and mammals. *Ecography* 28:403–416
- McCallum H, Barlow N, Hone J (2001) How should pathogen transmission be modelled? *Trends Ecol Evol* 16:295–300
- McLoughlin PD, Ferguson SH, Messier F (2000) Intraspecific variation in home range overlap with habitat quality: a comparison among brown bear populations. *Evol Ecol* 14:39–60
- Millsbaugh JJ, Brundige GC, Gitzen RA, Raedeke KJ (2004a) Herd organization of cow elk in Custer State Park, South Dakota. *Wildlife Soc B* 32:506–514
- Millsbaugh JJ, Gitzen RA, Kernohan BJ, Larson MA, Clay CL (2004b) Comparability of three analytical techniques to assess joint space use. *Wildlife Soc B* 32:148–157
- Mobæk R, Mysterud A, Loe LE, Holand Ø, Austrheim G (2012) Experimental evidence of density dependent activity pattern of a large herbivore in an alpine ecosystem. *Oikos* 121:364–369
- Morris DW (2003) Toward an ecological synthesis: a case for habitat selection. *Oecologia* 136:1–13
- Moyle K, Johnston D, Knott B, Melville-Smith R, Walker D (2009) Effect of stocking density on the growth, survival, and behavior of postpuerulus western rock lobster, *Panulirus cygnus* (George) (Decapoda: Palinuridae). *J World Aquac Soc* 40:255–265
- Nielsen CK, Woolf A (2001) Spatial organization of bobcats (*Lynx rufus*) in Southern Illinois. *Am Midl Nat* 146:43–52
- Nishi JS, Shury T, Elkin BT (2006) Wildlife reservoirs for bovine tuberculosis (*Mycobacterium bovis*) in Canada: strategies for management and research. *Vet Microbiol* 112:215–338
- Ostfeld RS, Lidicker WZ, Heske EJ (1985) The relationship between habitat heterogeneity, space use, and demography in a population of California voles. *Oikos* 45:433–442
- Pépin D, Gerard J-F (2008) Group dynamics and local population density dependence of group size in the Pyrenean chamois, *Rupicapra pyrenaica*. *Anim Behav* 75:361–369
- Pettit-Riley R, Estevez I, Russek-Cohen E (2002) Effects of crowding and access to perches on aggressive behaviour in broilers. *App Anim Behav Sci* 79:11–25
- Prange S, Jordan T, Hunter C, Gehrt SD (2006) New radiocollars for the detection of proximity among individuals. *Wildlife Soc B* 34:1333–1344

- R Development Core Team (2012) R: A Language and Environment for Statistical Computing. R Foundation for Statistical Computing, Vienna
- Ramsey D, Spencer N, Caley P, Efford M, Hansen K, Lam M, Cooper D (2002) The effects of reducing population density on contact rates between brushtail possums: implications for transmission of bovine tuberculosis. *J Appl Ecol* 39:806–818
- Rieucan G, Giraldeau L-A (2011) Exploring the costs and benefits of social information use: an appraisal of current experimental evidence. *Philos T R Soc Lond B* 366:949–957
- Robert K, Garant D, Pelletier F (2012) Keep in touch: does spatial overlap correlate with contact rate frequency? *J Wildlife Manage* 76:1670–1675
- Robert KR, Garant D, Vander Wal E, Pelletier F (2013) Context-dependent social behaviour: testing the interplay between season and kinship with raccoons. *J Zool* 290:199–207
- Rodgers AR, Carr AP, Smith L, Kie J (2005) Home Range Tools for ArcGIS 9.x. Ontario Ministry of Natural Resources. Centre for Northern Forest Ecosystem Research, Thunder Bay, Ontario
- Rowe JS (1972) Forest Regions of Canada. Information Canada, Ottawa, Ontario
- Ruckstuhl KE, Festa-Bianchet M (2001) Group choice by subadult bighorn rams: trade-offs between foraging efficiency and predator avoidance. *Ethology* 107:161–172
- Schauber EM, Storm DJ, Nielsen CK (2007) Effects of joint space use and group membership on contact rates among white-tailed deer. *J Wildlife Manage* 71:155–163
- Shury TK, Bergeson G (2011) Lesion distribution and epidemiology of *Mycobacterium bovis* in elk and white-tailed deer in south-western Manitoba, Canada. *Vet Med Int* 2011: Article ID 591980, 11 pages
- Steury TD, Murray DL (2003) Causes and consequences of individual variation in territory size in the American red squirrel. *Oikos* 101: 147–156
- Tompkins DM, Dunn AM, Smith MJ, Telfer S (2011) Wildlife diseases: from individuals to ecosystems. *J Anim Ecol* 80:19–38
- Vander Wal E (2011) Sex, friends, and disease: the social ecology of elk (*Cervus elaphus*) with implications for pathogen transmission. Dissertation, University of Saskatchewan
- Vander Wal E, McLoughlin PD, Brook RK (2011) Spatial and temporal factors influencing sightability of elk. *J Wildlife Manage* 75:1521–1526
- Vander Wal E, Yip H, McLoughlin PD (2012a) Sex-based differences in density-dependent sociality: an experiment with a gregarious ungulate. *Ecology* 93:206–212
- Vander Wal E, Paquet PC, Andrés JA (2012b) Influence of landscape and social interactions on transmission of disease in a social cervid. *Mol Ecol* 21:1271–1282
- Vander Wal E, van Beest FM, Brook RK (2013a) Density-dependent effects of group size are sex-specific in a gregarious ungulate. *PLoS ONE* 8:e53777
- Vander Wal E, Edye I, Paquet PC, Coltman DW, Bayne E, Brook RK, Andrés JA (2013b) Juxtaposition between host population structures: implications for pathogen transmission in a cervid community. *Evol Appl* 6:1001–1011
- Vander Wal E, Paquet PC, Messier F, McLoughlin PD (2013c) Effects of phenology and sex on social proximity in a gregarious ungulate. *Can J Zool* 91:601–609
- Walrath R, Van Deelen TR, VerCauteren KC (2011) Efficacy of proximity loggers for detection of contacts between maternal pairs of white-tailed deer. *Wildlife Soc B* 35:452–460
- Weckerly FW (1999) Social bonding and aggression in female Roosevelt elk. *Can J Zool* 77:1379–1384
- White TCL (2004) Limitation of populations by weather-driven changes in food: a challenge to density-dependent regulation. *Oikos* 105: 664–666
- White PCL, Harris S (1994) Encounters between red foxes (*Vulpes vulpes*): implications for territory maintenance, social cohesion and dispersal. *J Anim Ecol* 63:315–327
- Whitehead H (1997) Analysing animal social structure. *Anim Behav* 53: 1953–1067
- Whitehead H, Dufault S (1999) Techniques for analyzing vertebrate social structure using identified individuals: review and recommendations. *Adv Stud Behav* 28:33–74
- Wilson EO (1975) *Sociobiology: The New Synthesis*. Belknap, Cambridge, Massachusetts
- Wronski T, Apio A (2006) Home-range overlap, social vicinity and agonistic interactions denoting matrilineal organisation in bushbuck, *Tragelaphus scriptus*. *Behav Ecol Sociobiol* 59:819–828

### Author contributions

EVW conceived and designed the study. EVW collected the data. EVW and MPL analysed the data. EVW, MPL, and PDM interpreted the results. EVW, MPL, and PDM wrote and critically revised the manuscript.